

The Bretón-Manko equatorially antisymmetric binary configuration revisited

V. S. Manko[†], R. I. Rabadán[†] and E. Ruiz[‡]

[†]*Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN, A.P. 14-740, 07000 México D.F., Mexico*

[‡]*Instituto Universitario de Física Fundamental y Matemáticas,
Universidad de Salamanca, 37008 Salamanca, Spain*

The Bretón-Manko solution for two identical counter-rotating Kerr-Newman charged masses is rewritten in the physical parametrization involving Komar quantities. The new form of the solution in particular turns out to be very convenient for verifying a recent geometric inequality for interacting black holes with struts.

PACS numbers: 04.20.Jb, 04.70.Bw, 97.60.Lf

I. INTRODUCTION

The Bretón-Manko (BM) stationary axisymmetric electrovac solution of the Einstein-Maxwell equations [1] was constructed in 1995 with the aid of Sibgatullin's integral method [2, 3] and it describes a system of two identical counter-rotating Kerr-Newman (KN) sources, black holes or hyperextreme objects. The sources in the BM configuration are prevented from falling onto each other by a massless strut [4] which is removable only in the special case of equal masses and charges when the solution becomes a specialization of the Parker-Ruffini-Wilkins hyperextreme field [5] generated by means of the Perjés-Israel-Wilson method [6, 7]. The BM solution and its generalizations gave rise to a systematic study of equatorially antisymmetric spacetimes [8–10] which were shown to constitute a rather broad and interesting subfamily of stationary axisymmetric solutions. While the paper [1] mainly deals with the mathematical structure of BM solution and associated equilibrium problem, it would certainly be of interest to better explore the physical properties of the spacetime whose subextreme sector provides excellent opportunities for the study of two interacting KN black holes and for extension of the recent results on interacting Kerr sources [11, 12] to the case of non-vanishing electromagnetic field. It would be also desirable to use the general BM metric for testing the newly proposed geometric inequality [13] for multiple black holes separated by struts.

To make the BM solution more suitable for the physical analysis, in the present paper we shall rewrite it in a new parametrization involving the physical Komar quantities [14] and a separation distance as arbitrary parameters. This will enable us to demonstrate analytically that each KN black-hole constituent from the subextreme sector of the BM solution verifies the well-known Smarr's mass formula [15]. We will subsequently apply the BM solution to verifying a recent inequality on the bounds of the interaction force in black-hole systems with struts.

II. THE BM SOLUTION IN PHYSICAL PARAMETERS

The original BM solution arises from the axis data

$$\begin{aligned}\mathcal{E}(\rho=0, z) &= \frac{(z-k-M-ia')(z+k-M+ia')}{(z-k+M-ia')(z+k+M+ia')}, \\ \Phi(\rho=0, z) &= \frac{2Qz}{(z-k+M-ia')(z+k+M+ia')},\end{aligned}\tag{1}$$

representing the axis expressions of the Ernst complex potentials [16]. The parameters M , a' , Q and k in (1) are related, respectively, to the mass, angular momentum, charge of each KN source and to the separation distance, ρ and z are the Weyl-Papapetrou cylindrical coordinates. In the paper [1], Sibgatullin's method was applied to the data (1) for obtaining the corresponding Ernst potentials in the whole space, as well as the metric functions f , γ and ω entering the line element

$$ds^2 = f^{-1}[e^{2\gamma}(d\rho^2 + dz^2) + \rho^2 d\varphi^2] - f(dt - \omega d\varphi)^2.\tag{2}$$

The explicit original form of $\mathcal{E}$, Φ , f , γ and ω of the BM solution the reader can find in [1].

It should be pointed out that the parametrization used in [1] is not the most attractive one from the physical point of view because only the parameters M and Q coincide exactly with the physical mass and charge of each KN constituent; at the same time, the parameter a' does not coincide with the angular momentum per unit mass of any of the constituents, whereas the parameter k is not actually the coordinate distance between the centers of the constituents. In the absence of the electric charge Q (pure vacuum case) when the BK solution represents two

counter-rotating Kerr sources [17] a similar problem was already solved in the paper [11] by introducing the Komar angular momentum per unit mass a instead of a' , and the coordinate distance R between the centers of the sources instead of k (see Fig. 1).




FIG. 1: Location of identical counter-rotating KN black holes on the symmetry axis and the new parametrization.

Remarkably, transition from the parameters (a', k) to the physically more transparent parameters (a, R) can be performed in the general BM solution too, and after a considerable effort we have eventually been able to find the required reparametrized form of the data (1) and the form of an important quantity σ generalizing the analogous quantity in the vacuum case (cf. formula (3) of [11]):

$$\begin{aligned}\mathcal{E}(\rho=0, z) &= \frac{z^2 - 2Mz - (\frac{1}{2}MR + M^2 - \frac{1}{2}Q^2 - iMa)^2\mu}{z^2 + 2Mz - (\frac{1}{2}MR + M^2 - \frac{1}{2}Q^2 - iMa)^2\mu}, \\ \Phi(\rho=0, z) &= \frac{2Qz}{z^2 + 2Mz - (\frac{1}{2}MR + M^2 - \frac{1}{2}Q^2 - iMa)^2\mu},\end{aligned}\quad (3)$$

and

$$\sigma = \sqrt{M^2 - Q^2 - M^2a^2\mu}, \quad \mu := \frac{R^2 - 4M^2 + 4Q^2}{(MR + 2M^2 - Q^2)^2}. \quad (4)$$

The application of Sibgatullin's method to the data (3) yields the following new expressions for the Ernst potentials of the BM solution instead of the old ones:

$$\begin{aligned}\mathcal{E} &= \frac{A-B}{A+B}, \quad \Phi = \frac{C}{A+B}, \\ A &= (M^2 - Q^2)[4\sigma^2(R_+R_- + r_+r_-) + R^2(R_+r_+ + R_-r_-)] + [\sigma^2(R^2 - 4M^2 + 4Q^2) - M^2a^2R^2\mu] \\ &\quad \times (R_+r_- + R_-r_+) - 2iaMR\mu\sigma(MR + 2M^2 - Q^2)(R_+r_- - R_-r_+), \\ B &= 2MR\sigma\{R\sigma(R_+ + R_- + r_+ + r_-) - [2(M^2 - Q^2) + iMa\mu(MR + 2M^2 - Q^2)](R_+ - R_- - r_+ + r_-)\}, \\ C &= 2QR\sigma\{R\sigma(R_+ + R_- + r_+ + r_-) - [2(M^2 - Q^2) + iMa\mu(MR + 2M^2 - Q^2)](R_+ - R_- - r_+ + r_-)\} \\ &= QB/M, \\ R_{\pm} &= \sqrt{\rho^2 + (z + \frac{1}{2}R \pm \sigma)^2}, \quad r_{\pm} = \sqrt{\rho^2 + (z - \frac{1}{2}R \pm \sigma)^2}.\end{aligned}\quad (5)$$

On the other hand, the new expressions for the functions f , γ and ω of the BM solution have been found to have the form

$$\begin{aligned}
f &= \frac{AA^* - BB^* + CC^*}{(A+B)(A^*+B^*)}, \quad e^{2\gamma} = \frac{AA^* - BB^* + CC^*}{16R^4\sigma^4 R_+ R_- r_+ r_-}, \quad \omega = -\frac{\text{Im}[2G(A^* + B^*) + CI^*]}{AA^* - BB^* + CC^*}, \\
G &= -zB + R\sigma\{ (2M^2 - Q^2)[2\sigma(r_+ r_- - R_+ R_-) + R(R_- r_- - R_+ r_+)] \\
&\quad + M(R + 2\sigma)[R\sigma - 2(M^2 - Q^2) - iMa\mu(2M^2 - Q^2 + MR)](R_+ - r_-) \\
&\quad + M(R - 2\sigma)[R\sigma + 2(M^2 - Q^2) + iMa\mu(2M^2 - Q^2 + MR)](R_- - r_+) \}, \\
I &= -zC + QR\sigma\{ 2M[2\sigma(r_+ r_- - R_+ R_-) + R(R_- r_- - R_+ r_+)] \\
&\quad + (R + 2\sigma)[R\sigma - 2(M^2 - Q^2) - iMa\mu(2M^2 - Q^2 + MR)](R_+ - r_-) \\
&\quad + (R - 2\sigma)[R\sigma + 2(M^2 - Q^2) + iMa\mu(2M^2 - Q^2 + MR)](R_- - r_+) \}, \tag{6}
\end{aligned}$$

where an asterisk and $\text{Im}(x)$ denote complex conjugation and the imaginary part of x , respectively. Note that the metric function ω in (6) is presented in a simpler and more elegant form than in the paper [1].

For completeness, below we also write out the expressions defining the electric and magnetic potentials of the BM solution:

$$A_4 = \text{Re}\left(\frac{C}{A+B}\right), \quad A_3 = \text{Im}\left(\frac{I}{A+B}\right), \tag{7}$$

where $\text{Re}(x)$ means taking real part of x . The knowledge of the above A_4 and A_3 is needed in particular for the analysis of Smarr's mass formula.

III. SOME PHYSICAL PROPERTIES OF BM SOLUTION

The real valued σ determine a black-hole sector of BM solution, so in that case each KN constituent must verify Smarr's mass formula for black holes

$$M = \frac{1}{4\pi}\kappa S + 2\Omega^H J + \Phi^H Q, \tag{8}$$

which, apart from the mass M , angular momentum J and charge Q , contains four quantities to be evaluated on the horizon: the surface gravity κ , the area of the horizon S , horizon's angular velocity Ω^H and the electric scalar Φ^H . Since the constituents in BM solution only differ in the orientation of their angular momenta, it is sufficient to check formula (8) for solely one subextreme constituent, say, the upper one; the second constituent will then have opposite angular momentum $-J$ and horizon's velocity $-\Omega^H$ whose product leaves invariant the second term on the right hand side of (8). The known Komar characteristics associated with the upper constituent are M , $J = -Ma$ and Q , while the quantities κ , S , Ω^H and Φ^H can be calculated with the aid of the formulae [18, 19]

$$\kappa = \sqrt{-\omega^{-2}e^{-2\gamma}}, \quad S = \frac{4\pi\sigma}{\kappa}, \quad \Omega^H = \omega^{-1}, \quad \Phi^H = A_4 - \Omega^H A_3, \tag{9}$$

thus yielding in our case

$$\begin{aligned}
\kappa &= \frac{R\sigma}{\Delta}, \quad S = \frac{4\pi\Delta}{R}, \quad \Omega^H = -\frac{Ma\mu(MR + 2M^2 - Q^2)}{\Delta}, \quad \Phi^H = \frac{Q[(R + 2M)(M + \sigma) - 2Q^2]}{\Delta}, \\
\Delta &= 2M(R + 2M)(M + \sigma) - Q^2(R + 4M + 2\sigma). \tag{10}
\end{aligned}$$

A direct check shows that the above quantities satisfy identically the relation (8), therefore the subextreme constituents in the BM solution are indeed charged rotating black holes.

Mention that in the limit $R \rightarrow \infty$ (infinite separation) one recovers from (10) the characteristics of a single KN black hole [20], and at finite R , formulae (10) take into account the interaction between the black holes. The interaction force in the BM configuration is defined by a very simple expression (γ_0 is the value of the metric function γ on the strut)

$$\mathcal{F} = \frac{1}{4}(e^{-\gamma_0} - 1) = \frac{M^2 - Q^2}{R^2 - 4M^2 + 4Q^2}, \tag{11}$$

which coincides with the expression for the interaction force between two identical Reissner-Nordström non-rotating charged masses [21, 22] (cf. formula (30) of [23]).

The limiting case of two extreme counter-rotating black holes is defined by the condition $\sigma = 0$. Solving the latter for R , we find that M and Q must satisfy the inequality $M^2 > Q^2$ as a condition of the reality of R and non-vanishing a . On the other hand, solving $\sigma = 0$ for a , we get

$$a^2 = \frac{M^2 - Q^2}{M^2 \mu}, \quad (12)$$

and taking into account that $0 < M^2 \mu < 1$ for all $M > 0$, $R > 2\sqrt{M^2 - Q^2}$, we conclude that a black hole in the BM configuration needs a larger absolute value of the angular momentum in order to become extremal than a single Kerr black hole. The Ernst potentials and metric functions of the extreme BM solution have been worked out in explicit form in the paper [24].

Let us also comment, in connection with recent interest in geometrical inequalities for black holes (we refer the reader to a comprehensive review [25] on the topic), that the extreme BM solution provides the first example of a stationary axisymmetric electrovac spacetime for which the equality is achieved in a recent relation for black holes with struts derived by Gabach Clement [13]:

$$\sqrt{1 + 4\mathcal{F}} \geq \frac{\sqrt{(8\pi J)^2 + (4\pi Q^2)^2}}{S}. \quad (13)$$

Indeed, taking into account (12), the extremal values of S and $(8\pi J)^2 + (4\pi Q^2)^2$ are

$$\begin{aligned} S &= \frac{4\pi}{R} [(M^2 - Q^2)(R + 4M) + M^2 R], \\ (8\pi J)^2 + (4\pi Q^2)^2 &= \frac{16\pi^2 [(M^2 - Q^2)(R + 4M) + M^2 R]^2}{R^2 - 4M^2 + 4Q^2}, \end{aligned} \quad (14)$$

with which, and with formula (11), the equality in (13) can be easily verified. In general, the inequality (13) establishes lower bounds on the interaction force between the black-hole constituents in the BM solution.

We conclude this section by noticing that Smarr's formula gives rise to the inequality

$$M \geq 2\Omega^H J + \Phi^H Q \quad (15)$$

since the first term on the right hand side of (8) is a non-negative quantity. Then the equality in (15) will be achieved exclusively by the extremal black holes, while the sub-extreme black holes will verify a strict inequality.

IV. CONCLUSIONS

The new representation of BM solution elaborated in the present paper is remarkable in several aspects: it is the first and most simple example of a stationary axisymmetric electrovac spacetime entirely written in physical parameters and able to describe the field of two interacting KN black holes (previous results on the physical parametrization of binary configurations were obtained either for a vacuum [11] or static electrovac [23] cases). It considerably simplifies the analysis of physical properties of the BM solution and permits one to get concise expressions for the basic characteristics of interacting KN black holes and use them in concrete applications. It also motivates the search for physical parametrizations of more general configurations of KN black holes than the BM one, for instance by introducing appropriately the charges into the vacuum solution for non-equal counter-rotating Kerr black holes [26]. Last but not least, an interesting aspect of the reparametrized BM solution is connected with a recent discovery by Dain and Ortiz [27] of the correspondence between stationary axisymmetric spacetimes with a strut and non-stationary axisymmetric spacetimes via the notion of momentary stationary data, thus suggesting that the BM solution could in principle shed some new light on the non-radiating binary non-stationary systems too.

Acknowledgements

We are thankful to Erasmo Gómez for technical computer support during the involved analytic calculations. This work was partially supported by CONACYT, Mexico, and by Ministerio de Ciencia y Tecnología, Spain, under the

Project FIS2009-07238.

-
- [1] N. Bretón and V. S. Manko, *Class. Quantum Grav.* **12** (1995) 1969.
 - [2] N. R. Sibgatullin, *Oscillations and Waves in Strong Gravitational and Electromagnetic Fields* (Springer-Verlag, Berlin, 1991).
 - [3] V. S. Manko and N. R. Sibgatullin, *Class. Quantum Grav.* **10** (1993) 1383.
 - [4] W. Israel, *Phys. Rev. D* **15** (1977) 935.
 - [5] L. Parker, R. Ruffini and D. Wilkins, *Phys. Rev. D* **7** (1973) 2874.
 - [6] Z. Perjés, *Phys. Rev. Lett.* **27** (1971) 1668.
 - [7] W. Israel and G. A. Wilson, *J. Math. Phys.* **13** (1972) 865.
 - [8] F. J. Ernst, V. S. Manko and E. Ruiz, *Class. Quantum Grav.* **23** (2006) 4945.
 - [9] F. J. Ernst, V. S. Manko and E. Ruiz, *Class. Quantum Grav.* **24** (2007) 2193.
 - [10] J. Sod-Hoffs and E. D. Rodchenko, *Class. Quantum Grav.* **24** (2007) 4617.
 - [11] V. S. Manko, E. D. Rodchenko, E. Ruiz and B. I. Sadovnikov, *Phys. Rev. D* **78** (2008) 124014.
 - [12] C. A. R. Herdeiro and C. Rebelo, *J. High Energy Phys.* **10** (2008) 017.
 - [13] M. E. Gabach Clement, *Class. Quantum Grav.* **29** (2012) 165008.
 - [14] A. Komar, *Phys. Rev.* **113** (1959) 934.
 - [15] L. Smarr, *Phys. Rev. Lett.* **30** (1973) 71.
 - [16] F. J. Ernst, *Phys. Rev.* **168** (1968) 1415.
 - [17] R. P. Kerr, *Phys. Rev. Lett.* **11** (1963) 237.
 - [18] B. Carter, in *General Relativity, an Einstein Centenary Survey*, ed. S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1979), p. 294.
 - [19] A. Tomimatsu, *Prog. Theor. Phys.* **72** (1984) 73.
 - [20] E. T. Newman, E. Couch, K. Chinnapared, A. Exton, A. Prakash and R. Torrence, *J. Math. Phys.* **6** (1965) 918.
 - [21] H. Reissner, *Ann. Physik* **50** (1916) 106.
 - [22] G. Nordström, *Proc. Kon. Ned. Akad. Wet.* **20** (1918) 1238.
 - [23] V. S. Manko, *Phys. Rev. D* **76** (2007) 124032.
 - [24] V. S. Manko, E. Ruiz and M. B. Sadovnikova, *Phys. Rev. D* **84** (2011) 064005.
 - [25] S. Dain, *Class. Quantum Grav.* **29** (2012) 073001.
 - [26] I. Cabrera-Munguia, C. Lämmerzahl and A. Macías, arXiv:1302.4843 [gr-qc].
 - [27] S. Dain and O. E. Ortiz, *Phys. Rev. D* **80** (2009) 024045.